



Product Datasheet

Product Name	Prolactin Soluble Receptor Human Recombinant
Cata No	CB501329
Source	<i>Escherichia Coli.</i>
Synonyms	PRL-R, hPRLrI.

Description

Prolactin is a pituitary hormone that plays a role in the stimulation of milk production, salt and water regulation, growth, development and reproduction. The primary step in its action is the binding to a specific membrane receptor (prolactin receptor) which belongs to the superfamily of class 1 cytokine receptors. Prolactin is a hormone involved in a range of significant functions including ion transport and osmoregulation, stimulation of milk, protein synthesis as well as the regulation of numerous reproductive functions. Prolactin exerts its influence on different cell types through a signal transduction pathway which begins with the binding of the hormone to a transmembrane Prolactin receptor. PRLR varies in size (short and long forms) with tissue source and species, from ~40 kDa to 100 kDa. The PRL-R consists of at least 3 separate domains: an extracellular region with 5 cysteines which contains the prolactin binding site, a single transmembrane domain and a cytoplasmic region, the length of which appears to influence ligand binding and regulate cellular function. Extra Cellular Domain Prolactin Receptor produced in E.Coli is a non-glycosylated, Polypeptide chain Human Recombinant containing 210 amino acids and having a molecular mass of 23.97 kDa. The Prolactin Receptor is purified by proprietary chromatographic techniques according to Bignon et al. (1994) JBC 269; 3318-24 and tested according to Gertler et al. (1996) JBC 271; 24482-91.

Physical Appearance

Sterile filtered white lyophilized powder.

Biological Activity

Activity is determined by the dose-dependant inhibition of Prolactin stimulated proliferation of Nb2 cells and by high affinity binding of ovine Prolactin and other lactogenic hormones in 1:1 molar ratio

Purity

Greater than 97.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.
- (c) Gel filtration at pH 8 under non denaturative conditions.

Formulation

The Prolactin Receptor was lyophilized from a concentrated (0.4mg/ml) solution with 0.0045mM NaHCO₃.

Reconstitution

It is recommended to reconstitute the lyophilized PRLR in sterile 18M-cm H₂O not less than 100µg/ml and not more than 1 mg/ml, which can then be further diluted to other aqueous solutions.

Stability

Lyophilized PRL-R although stable at room temperature for 1-2 weeks, should be stored desiccated below -18°C or preferably even at -80°C to prevent dimer formation. Upon reconstitution PRL-R should be stored sterile at 4°C between 2-7 days and for future use below -18°C. For long term

*** For Non-Clinical Research Use Only ***



California Bioscience

83103 Avenue 48, Ste.1B #204
Coachella, CA 92236 USA
Phone : +1.6268339877
Email : info@cali-bio.com

storage at 4°C it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles as they cause oligomerization of the protein.

Sequence

The sequence of the first product was determined to be Ala-Gly-Lys-Pro-Glu-Ile.

Product Data Sheet

*** For Non-Clinical Research Use Only ***